

BOUNDARY SLOPES AND THE LOGARITHMIC LIMIT SET

STEPHAN TILLMANN

ABSTRACT. The A -polynomial of a manifold whose boundary consists of a single torus is generalised to an *eigenvalue variety* of a manifold whose boundary consists of a finite number of tori, and the set of strongly detected boundary curves is determined by Bergman's *logarithmic limit set*, which describes the exponential behaviour of the eigenvalue variety at infinity. This enables one to read off the detected boundary curves of a multi-cusped manifold in a similar way to the 1-cusped case, where the slopes are encoded in the Newton polygon of the A -polynomial.

INTRODUCTION

Let V be an algebraic variety in $(\mathbb{C} - \{0\})^m$. The first section discusses the *logarithmic limit set* V_∞ of V , which we use to define the set of ideal points of V . We show that each point with rational coordinate ratios in V_∞ is an ideal point of a curve in V . Thus, the logarithmic limit set lends itself to applications of Culler–Shalen theory, where essential surfaces are associated to ideal points of curves in the character variety of a 3-manifold.

Given a compact, orientable, irreducible 3-manifold with boundary consisting of a single torus, one calls boundary slopes of essential surfaces associated to ideal points of the character variety *strongly detected*. It is shown in [5] that the slope of each side of the Newton polygon of the A -polynomial is a strongly detected boundary slope.

In the second section, we generalise the A -polynomial to define an *eigenvalue variety* of a manifold with boundary consisting of h tori. We describe the *strongly detected boundary curves*, i.e. the boundary curves of essential surfaces associated to ideal points of the character variety, via a map from the logarithmic limit set of the eigenvalue variety to the $(2h - 1)$ -sphere.

The third section contains a computation of the A -polynomials of torus knots. These seem to be the only infinite families of A -polynomials currently known.

Date: February 1, 2008.

1. BERGMAN'S LOGARITHMIC LIMIT SET

1.1. Logarithmic limit set. Let V be a subvariety of $(\mathbb{C} - \{0\})^m$ defined by an ideal J , and let $\mathbb{C}[X^\pm] = \mathbb{C}[X_1^{\pm 1}, \dots, X_m^{\pm 1}]$. Bergman gives the following three descriptions of a *logarithmic limit set* in [1]:

1. Define $V_\infty^{(a)}$ as the set of limit points on S^{m-1} of the set of real m -tuples in the interior of B^m :

$$(1) \quad \left\{ \frac{(\log |x_1|, \dots, \log |x_m|)}{\sqrt{1 + \sum (\log |x_i|)^2}} \mid x \in V \right\}.$$

2. Define $V_\infty^{(b)}$ as the set of m -tuples

$$(2) \quad (-v(X_1), \dots, -v(X_m))$$

as v runs over all real-valued valuations on $\mathbb{C}[X^\pm]/J$ satisfying $\sum v(X_i)^2 = 1$.

3. Define the support $s(f)$ of an element $f \in \mathbb{C}[X^\pm]$ to be the set of points $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{Z}^m$ such that $X^\alpha = X_1^{\alpha_1} \cdots X_m^{\alpha_m}$ occurs with non-zero coefficient in f . Then define $V_\infty^{(c)}$ to be the set of $\xi \in S^{m-1}$ such that for all non-zero $f \in J$, the maximum value of the dot product $\xi \cdot \alpha$ as α runs over $s(f)$ is assumed at least twice.

Bergman shows in [1] that the second and third descriptions are equivalent, and it follows from Bieri and Groves [2] that all descriptions are equivalent:


Theorem 1 (Bergman, Bieri–Groves). $V_\infty^{(a)} = V_\infty^{(b)} = V_\infty^{(c)}$.

We therefore write V_∞ for the logarithmic limit set of the variety V . A generic picture for the projection of a variety in $(\mathbb{C} - \{0\})^2$ into B^2 and its logarithmic limit set in S^1 is given in Figure 1. The ideal generated by a set of polynomials $\{f_i\}_i$ shall be denoted by $I(f_i)_i$, and the variety generated by the ideal I is denoted by $V(I)$ or $V(f_i)_i$, if $I = I(f_i)_i$.

1.2. Newton Polytopes. The third description of the logarithmic limit set can be illustrated using spherical duals of Newton polytopes. Let $f \in \mathbb{C}[X^\pm]$ be given as an expression $f = \sum_{\alpha \in \mathbb{Z}^m} a_\alpha X^\alpha$. Then the *Newton polytope* of f is the convex hull of $s(f) = \{\alpha \in \mathbb{Z}^m \mid a_\alpha \neq 0\}$. Thus, if $s(f) = \{\alpha_1, \dots, \alpha_k\}$, then

$$(3) \quad \text{Newt}(f) = \text{Conv}(s(f)) = \left\{ \sum_{i=1}^k \lambda_i \alpha_i \mid \lambda_i \geq 0, \sum_{i=1}^k \lambda_i = 1 \right\}.$$

Note that $\text{Newt}(0) = \emptyset$, and since $V(0) = (\mathbb{C} - \{0\})^m$, we have $V(0)_\infty = S^{m-1}$.


 FIGURE 1. A logarithmic limit set in S^1

For arbitrary subsets A, B of $\mathbb{R}^m$, let $A + B = \{a + b \mid a \in A, b \in B\}$. The following facts about Newton polytopes of polynomials are well known (cf. [7]):

- (1) Let $f, g \in \mathbb{C}[X^\pm]$ such that $fg \neq 0$. Then $\text{Newt}(fg) = \text{Newt}(f) + \text{Newt}(g)$.
- (2) $\text{Newt}(f + g) \subseteq \text{Conv}(s(f) \cup s(g)) = \text{Conv}(\text{Newt}(f) \cup \text{Newt}(g))$.

1.3. Spherical Duals. The *spherical dual* of a bounded convex polytope P in $\mathbb{R}^m$ is the set of vectors ξ of length 1 such that the supremum $\sup_{\alpha \in P} \alpha \cdot \xi$ is achieved for more than one α , and it is denoted by $\text{Sph}(P)$. Geometrically, $\text{Sph}(f)$ consists of all outward pointing unit normal vectors to the support planes of P which meet P in more than one point. If P is the Newton polytope of a non-zero polynomial f , then the spherical dual of $\text{Newt}(f)$ is also denoted by $\text{Sph}(f)$. The following lemma is immediate from the third description of the logarithmic limit set:

Lemma 2. *Let V be an algebraic variety in $(\mathbb{C} - \{0\})^m$ defined by the ideal J . Then V_∞ is the intersection over all non-zero elements of J of the spherical duals of their Newton polytopes.*

Spherical duals are easy to visualise in low dimensions. Given a convex polygon P in $\mathbb{R}^2$, its spherical dual is the collection of points on the unit circle defined by outward pointing unit normal vectors to edges of P . Given a convex polyhedron in $\mathbb{R}^3$, we obtain the vertices of its spherical dual again as points on the unit sphere S^2 arising from outward pointing unit normal vectors to faces. We then join two of these points along the shorter geodesic arc in S^2 if the corresponding faces have a common edge. This gives a finite graph in S^2 .

1.4. If the ideal J is principal, defined by an element $f \neq 0$, then Bergman states in [1] that the set $V(f)_\infty$ is precisely $Sph(f)$. We give a proof using the following lemma.

Lemma 3. *Let $f, g \in \mathbb{C}[X^\pm]$. If $fg \neq 0$, then $Sph(fg) = Sph(f) \cup Sph(g)$.*

Proof. If $\xi \in Sph(f)$, then there are distinct $\alpha_0, \alpha_1 \in Newt(f)$ such that

$$\alpha_0 \cdot \xi = \alpha_1 \cdot \xi = \sup_{\alpha \in Newt(f)} \alpha \cdot \xi.$$

Since $Newt(g)$ is convex and bounded, there is $\beta_0 \in Newt(g)$ such that

$$\beta_0 \cdot \xi = \sup_{\beta \in Newt(g)} \beta \cdot \xi.$$

Since $Newt(fg) = Newt(f) + Newt(g)$, it follows that $\beta_0 + \alpha_0, \beta_0 + \alpha_1 \in Newt(fg)$ and

$$(\alpha_0 + \beta_0) \cdot \xi = (\alpha_1 + \beta_0) \cdot \xi = \sup_{\gamma \in Newt(fg)} \gamma \cdot \xi.$$

Thus, $Sph(f) \subseteq Sph(fg)$, and by symmetry $Sph(g) \subseteq Sph(fg)$.

Now then assume that $\xi \in Sph(fg)$. Then there are distinct γ_0 and γ_1 in $Newt(fg)$ such that

$$\gamma_0 \cdot \xi = \gamma_1 \cdot \xi = \sup_{\gamma \in Newt(fg)} \gamma \cdot \xi.$$

Furthermore, there are $\alpha_i \in Newt(f)$ and $\beta_i \in Newt(g)$ such that $\gamma_i = \alpha_i + \beta_i$. If $\alpha_0 \cdot \xi < \alpha_1 \cdot \xi$, then $\gamma_0 \cdot \xi < (\alpha_1 + \beta_0) \cdot \xi$ and $\alpha_1 + \beta_0 \in Newt(fg)$, contradicting the choice of γ_0 . Thus, $\alpha_0 \cdot \xi = \alpha_1 \cdot \xi$, and similarly $\beta_0 \cdot \xi = \beta_1 \cdot \xi$. A similar argument shows:

$$\alpha_0 \cdot \xi = \alpha_1 \cdot \xi = \sup_{\alpha \in Newt(f)} \alpha \cdot \xi,$$

and similarly for the β_i . Since $\gamma_0 \neq \gamma_1$, either $\alpha_0 \neq \alpha_1$ or $\beta_0 \neq \beta_1$, and hence either $\xi \in Sph(f)$ or $\xi \in Sph(g)$. ■

Proposition 4. *Let $f \in \mathbb{C}[X^\pm]$. If $f \neq 0$, then $V(f)_\infty = Sph(f)$.*

Proof. Since each non-zero element of $I(f)$ is of the form fg for some non-zero $g \in \mathbb{C}[X^\pm]$, Lemma 2 and Lemma 3 yield:

$$V(f)_\infty = \bigcap_{0 \neq g \in \mathbb{C}[X^\pm]} Sph(fg) = \bigcap_{0 \neq g \in \mathbb{C}[X^\pm]} (Sph(f) \cup Sph(g)) = Sph(f),$$

since $Sph(1) = \emptyset$. ■

Logarithmic limit sets of varieties defined by a principal ideal are therefore completely determined by a generator of the ideal, and readily computable. If the ideal J is not principal, then $V(J)_\infty$ is the intersection of the spherical duals of the Newton polytopes of *finitely* many elements in J . This follows from Bieri and Groves [2]. However, there is presently no known algorithm to determine a suitable finite set of elements of J . Conjecturally, a Gröbner basis may suffice.

1.5. As Bergman notes, the spherical dual of the convex hull of a finite subset $A \subset \mathbb{Z}^m$ of cardinality r is a finite union of convex spherical polytopes. It is the union over all $\alpha_0, \alpha_1 \in A$ of the set of ξ satisfying the $2r$ inequalities

$$\alpha_0 \cdot \xi \geq \alpha \cdot \xi \quad \text{and} \quad \alpha_1 \cdot \xi \geq \alpha \cdot \xi,$$


where α ranges over A . This may be useful for calculations, but it also shows that corners of the convex spherical polytopes which we consider have rational coordinate ratios, and points with rational coordinate ratios are dense in the polytope. Such a polytope is called a *rational convex spherical polytope*. We summarise further results by Bergman and Bieri–Groves as follows:

Theorem 5 (Bergman, Bieri–Groves). *The logarithmic limit set V_∞ of an algebraic variety V in $(\mathbb{C} - \{0\})^m$ is a finite union of rational convex spherical polytopes. The maximal dimension for a polytope in this union is achieved by at least one member P in this union, and we have*

$$\dim_{\mathbb{R}} V_\infty = \dim_{\mathbb{R}} P = (\dim_{\mathbb{C}} V) - 1.$$

1.6. **Whitehead link.** Bergman asks in [1] whether every polytope in the above union always has the same dimension. In [18] it is shown that the irreducible component in the $PSL_2(\mathbb{C})$ -eigenvalue variety of the Whitehead link complement $\mathcal{W}$ which arises from the Dehn surgery component has logarithmic limit set consisting of five components, four of which are points, and one of which is a union of geodesic arcs. A schematic picture is given in Figure 2.

It would be interesting to modify Bergman’s question and ask whether each connected component of the logarithmic limit set of an irreducible variety can always be expressed as a union of polytopes all having the same dimension when $\dim_{\mathbb{C}} V > 2$.

FIGURE 2. The logarithmic limit set of $\overline{\mathfrak{E}}_0(\mathcal{W})$

1.7. Curve finding lemma. We conclude this section with a lemma, which is useful for applications of Culler–Shalen theory.

Lemma 6. *Let V be an algebraic variety in $(\mathbb{C} - \{0\})^m$, and $\xi \in V_\infty$ be a point with rational coordinate ratios. Then there is a curve C , i.e. an irreducible subvariety of complex dimension one, in V such that $\xi \in C_\infty$.*

Proof. By a lemma of Bergman ([1], p.464), the point ξ is contained in the logarithmic limit set of an irreducible component of V . Thus, we may assume that V is irreducible for the purpose of this proof.

Assume that V is defined by the ideal I in $\mathbb{C}[X^\pm] = \mathbb{C}[X_1^{\pm 1}, \dots, X_m^{\pm 1}]$. We follow word by word Bergman’s construction starting after “(4)” on page 465, up to the sentence “Hence the points $u_y \log |y| \in B^m$ will approach $(0, \dots, 0, -1)$.” on page 466. Here u_y denotes $(\sqrt{1 + \sum (\log(|y_i|))^2})^{-1}$. Our adaption reads as follows:

Let ξ be a point of V_∞ the ratios of whose coordinate functions are rational. Making use of the action of $GL_m(\mathbb{Z})$ on $\mathbb{C}[X^\pm]$ and the induced action on S^{m-1} (not isometric!), we can reduce to the case where $\xi = (0, \dots, 0, -1)$.

Let R' designate the ring $\mathbb{C}[X_1^{\pm 1}, \dots, X_{m-1}^{\pm 1}; X_m] \subset R = \mathbb{C}[X^\pm]$, and R_0 the ring $\mathbb{C}[X_1^{\pm 1}, \dots, X_{m-1}^{\pm 1}]$, into which we map R' by sending X_m to 0. Let I' designate $I \cap R'$, and I_0 the image of I' in R_0 . Thus, I_0 consists of the components of degree 0 in X_m of elements of I' . Let V' designate the subvariety of $(\mathbb{C} - \{0\})^{m-1} \times \mathbb{C}$ defined by I' , and $V_0 = V' \cap (\mathbb{C} - \{0\})^{m-1} \times \{0\}$, the subvariety of $(\mathbb{C} - \{0\})^{m-1} \times \{0\}$ defined by I_0 . V' will be the Zariski closure of V in $(\mathbb{C} - \{0\})^{m-1} \times \mathbb{C}$, and $V = V' - V_0$.

If $(0, \dots, 0, -1) \in V_\infty$, this means that every nonzero $a \in I$ has more than one term of minimal degree in X_m . Hence every nonzero element of I_0 has more than one term, i.e. I_0 is a proper ideal of R_0 . Then V_0 is non-empty: let $x \in V_0$.

Now V will in fact be dense in V' under the topology induced by the absolute value on $\mathbb{C}$. [...] Hence we can obtain x as a limit of points $y \in V$. As y approaches x , $\log |y_i|$ will approach the finite values $\log |x_i|$ for $i = 1, \dots, m-1$, but $\log |y_m|$ will approach $-\infty$. Hence the points $u_y \log |y| \in B^m$ will approach $(0, \dots, 0, -1)$.

We construct a curve C in V such that $\xi = (0, \dots, 0, -1) \in C_\infty$ using the following two facts from [13]:

- (1) (1.14) Let Y be an irreducible affine variety. If X is a proper subvariety of Y , then $\dim X < \dim Y$.
- (2) (3.14) If X is an r -dimensional affine variety and $f_1, \dots, f_k$ are polynomial functions on X , then every component of $X \cap V(f_1, \dots, f_k)$ has dimension $\geq r - k$.

Since V_0 is a proper subvariety of V' , obtained by intersection with a hyperplane, we have $\dim V_0 = \dim V' - 1$, and since $V = V' - V_0$, we have $\dim V = \dim V'$. We need to find a curve C' in V' which has non-trivial intersection with both V_0 and V , since then $C = C' \cap V$ is a curve in V such that $\xi \in C_\infty$.

Let $\dim V = d$. Let $x \in V_0$ and choose a polynomial f_1 on V_0 such that $f_1(x) = 0$, but f_1 is not identical to 0 on V_0 . Then f_1 is not identical to 0 on V' and each component of the variety $V' \cap V(f_1)$ has dimension exactly $d - 1$, and intersects V_0 in a $(d - 2)$ -dimensional subvariety. Let V_1 be a component of $V' \cap V(f_1)$ containing x .

Choose a polynomial f_2 such that $f_2(x) = 0$, but f_2 is not identical to 0 on $V_1 \cap V_0$. Then each component of $V_1 \cap V(f_2) = V' \cap V(f_1, f_2)$ has dimension exactly $d - 2$ and intersects V_0 in a $(d - 3)$ -dimensional subvariety. Let V_2 be a component of $V' \cap V(f_1, f_2)$ containing x .

We now proceed inductively for $k = 2, \dots, d - 2$. Given a component V_k of $V' \cap V(f_1, \dots, f_k)$ which is a $(d - k)$ -dimensional subvariety of V' , with the property that it contains x , but also points of $V' - V_0$, we let f_{k+1} be a polynomial function on V_k which vanishes on x but not on all of $V_k \cap V_0$. Then each component of $V' \cap V(f_1, \dots, f_{k+1})$ has dimension exactly $d - k - 1$ and intersects V_0 in a $(d - k - 2)$ -dimensional subvariety, and we let V_{k+1} be a component of $V' \cap V(f_1, \dots, f_{k+1})$ containing x .

Then $V_{d-1} \cap V$ is a curve in V with the desired property. ■

2. THE EIGENVALUE VARIETY

2.1. Essential surfaces. A *surface* S in a compact 3-manifold M will always mean a 2-dimensional PL submanifold *properly embedded* in M , that is, a closed subset of M with $\partial S = S \cap \partial M$. If M is not compact, we replace it by a compact core. An embedded sphere S^2 in a 3-manifold M is called *incompressible* if it does not bound an embedded ball in M , and a manifold is *irreducible* if it contains no incompressible 2-spheres.

An orientable surface S without 2-sphere or disc components in an orientable 3-manifold M is called *irreducible* if for each disc $D \subset M$ with $D \cap S = \partial D$ there is a disc $D' \subset S$ with $\partial D' = \partial D$. We will also use the following definition:

Definition 7. [15] *A surface S in a compact, irreducible, orientable 3-manifold is said to be essential if it has the following five properties:*

- (1) *S is bicollared;*
- (2) *the inclusion homomorphism $\pi_1(S_i) \rightarrow \pi_1(M)$ is injective for every component S_i of S ;*
- (3) *no component of S is a 2-sphere;*
- (4) *no component of S is boundary parallel;*
- (5) *S is nonempty.*

2.2. ∂ -incompressible. The following definition and facts can be found in [11]. A (properly embedded) surface S in a compact 3-manifold M is *∂ -incompressible* if for each disc $D \subset M$ such that ∂D splits into two arcs α and β meeting only at their common endpoints with $D \cap S = \alpha$ and $D \cap \partial M = \beta$ there is a disc $D' \subset S$ with $\alpha \subset \partial D'$ and $\partial D' - \alpha \subset \partial S$. A surface S is incompressible and ∂ -incompressible if and only if each component of S is incompressible and ∂ -incompressible.

Lemma 8. [11] *Let S be a connected incompressible surface in the irreducible 3-manifold M , with ∂S contained in torus boundary components of M .*

Then either S is ∂ -incompressible or S is a boundary parallel annulus.

For the remainder of this section, we let M be a compact, orientable, irreducible 3-manifold with non-empty boundary consisting of a disjoint union of h tori. It follows from Definition 7 and Lemma 8 that an essential surface in M is ∂ -incompressible.

2.3. Boundary curve space. Denote the boundary tori of M by $T_1, \dots, T_h$, and choose an oriented meridian $\mathcal{M}_i$ and an oriented longitude l_i for each T_i . Let S be an incompressible

and ∂ -incompressible surface with non-empty boundary in M . There are coprime integers p_i, q_i such that S meets T_i in n_i curves parallel to $p_i\mathcal{M}_i + q_i l_i$. We thus obtain a point

$$(n_1 p_1, n_1 q_1, \dots, n_h p_h, n_h q_h) \in \mathbb{Z}^{2h} - \{0\}.$$

We view the above point as an element in $\mathbb{R}P^{2h-1}$ since most methods of detecting essential surfaces only do so up to projectivisation. Furthermore, we ignore the orientation of ∂S , so the loop $p_i\mathcal{M}_i + q_i l_i$ is equivalent to $-p_i\mathcal{M}_i - q_i l_i$. All we have to know are the relative signs of p_i and q_i . This equivalence induces a $\mathbb{Z}_2^{h-1}$ action on $\mathbb{R}P^{2h-1}$, and the closure of the set of *projectivised boundary curve coordinates*:

$$[n_1 p_1, n_1 q_1, \dots, n_h p_h, n_h q_h] \in \mathbb{R}P^{2h-1} / \mathbb{Z}_2^{h-1}$$

arising from incompressible and ∂ -incompressible surfaces with non-empty boundary in M is called the *boundary curve space* of M and denoted by $\mathfrak{BC}(M)$. Note that $\mathbb{R}P^{2h-1} / \mathbb{Z}_2^{h-1} \cong S^{2h-1}$ (see [10]). We will use the following result.

Theorem 9. [10] *Let M be a compact, orientable, irreducible 3-manifold with non-empty boundary consisting of a disjoint union of h tori. Then $\dim_{\mathbb{R}} \mathfrak{BC}(M) < h$.*

2.4. Character variety. Let Γ be a finitely generated group. The set of representations of Γ into $SL_2(\mathbb{C})$ is $\mathfrak{R}(\Gamma) = \text{Hom}(\Gamma, SL_2(\mathbb{C}))$. This set is called the *representation variety* of Γ , and regarded as an affine algebraic set. Each $\gamma \in \Gamma$ defines a *trace function* $I_\gamma : \mathfrak{R}(\Gamma) \rightarrow \mathbb{C}$ by $I_\gamma(\rho) = \text{tr } \rho(\gamma)$, which is an element of the coordinate ring $\mathbb{C}[\mathfrak{R}(\Gamma)]$.

Two representations are *equivalent* if they differ by an inner automorphism of $SL_2(\mathbb{C})$. A representation is *irreducible* if the only subspaces of $\mathbb{C}^2$ invariant under its image are trivial. This is equivalent to saying that the representation cannot be conjugated to a representation by upper triangular matrices. Otherwise a representation is *reducible*.

For each $\rho \in \mathfrak{R}(\Gamma)$, its *character* is the function $\chi_\rho : \Gamma \rightarrow \mathbb{C}$ defined by $\chi_\rho(\gamma) = \text{tr } \rho(\gamma)$. It turns out that irreducible representations are determined by characters up to equivalence, and that the reducible representations form a closed subset $\mathfrak{Red}(\Gamma)$ of $\mathfrak{R}(\Gamma)$.

The collection of characters $\mathfrak{X}(\Gamma)$ can be regarded as an affine algebraic set, which is called the *character variety*. There is a regular map $t : \mathfrak{R}(\Gamma) \rightarrow \mathfrak{X}(\Gamma)$ taking representations to characters.

If Γ is the fundamental group of a topological space M , we also write $\mathfrak{R}(M)$ and $\mathfrak{X}(M)$ instead of $\mathfrak{R}(\Gamma)$ and $\mathfrak{X}(\Gamma)$ respectively.

There also is a character variety arising from representations into $PSL_2(\mathbb{C})$, see Boyer and Zhang [3], and the relevant objects are denoted by placing a bar over the previous notation. The natural map $q : \mathfrak{X}(\Gamma) \rightarrow \overline{\mathfrak{X}}(\Gamma)$ is finite-to-one, but in general not onto. As with the $SL_2(\mathbb{C})$ -character variety, there is a surjective “quotient” map $\bar{t} : \overline{\mathfrak{R}}(\Gamma) \rightarrow \overline{\mathfrak{X}}(\Gamma)$, which is constant on conjugacy classes, and with the property that if $\bar{\rho}$ is an irreducible representation, then $\bar{t}^{-1}(\bar{t}(\bar{\rho}))$ is the orbit of $\bar{\rho}$ under conjugation. Again, a representation is irreducible if it is not conjugate to a representation by upper triangular matrices.

2.5. Dehn surgery component. If a 3-manifold M admits a complete hyperbolic structure of finite volume, then there is a discrete and faithful representation $\pi_1(M) \rightarrow PSL_2(\mathbb{C})$. This representation is necessarily irreducible, as hyperbolic geometry otherwise implies that M has infinite volume. If M is not compact, then a *compact core* of M is a compact manifold $\overline{M}$ such that M is homeomorphic to the interior of $\overline{M}$. We will rely on the following result:

Theorem 10 (Thurston). [16, 15] *Let M be a complete hyperbolic 3-manifold of finite volume with h cusps, and let $\bar{\rho}_0 : \pi_1(M) \rightarrow PSL_2(\mathbb{C})$ be a discrete and faithful representation associated to the complete hyperbolic structure. Then $\bar{\rho}_0$ admits a lift ρ_0 into $SL_2(\mathbb{C})$ which is still discrete and faithful. The (unique) irreducible component $\mathfrak{X}_0$ in the $SL_2(\mathbb{C})$ -character variety containing the character χ_0 of ρ_0 has (complex) dimension h .*

Furthermore, if $T_1, \dots, T_h$ are the boundary tori of a compact core of M , and if γ_i is a non-trivial element in $\pi_1(M)$ which is carried by T_i , then $\chi_0(\gamma_i) = \pm 2$ and χ_0 is an isolated point of the set

$$X^* = \{\chi \in \mathfrak{X}_0 \mid I_{\gamma_1}^2 = \dots = I_{\gamma_h}^2 = 4\}.$$

The respective irreducible components containing the so-called *complete representations* ρ_0 and $\bar{\rho}_0$ are denoted by $\mathfrak{R}_0(M)$ and $\overline{\mathfrak{R}}_0(M)$ respectively. In particular, $t(\mathfrak{R}_0) = \mathfrak{X}_0$ and $\bar{t}(\overline{\mathfrak{R}}_0) = \overline{\mathfrak{X}}_0$ are called the respective *Dehn surgery components* of the character varieties of M , since the holonomy representations of hyperbolic manifolds or orbifolds obtained by performing high order Dehn surgeries on M are near $\bar{\rho}_0$ (see [16]).

2.6. Eigenvalue variety. Given a presentation of $\pi_1(M)$ with a finite number, n , of generators, $\gamma_1, \dots, \gamma_n$, we introduce four affine coordinates (representing matrix entries) for each generator, which we denote by g_{ij} for $i = 1, \dots, n$ and $j = 1, 2, 3, 4$. We view

$\mathfrak{R}(M)$ as a subvariety of $\mathbb{C}^{4n}$ defined by an ideal J in $\mathbb{C}[g_{11}, \dots, g_{n4}]$. There are elements $I_\gamma \in \mathbb{C}[g_{11}, \dots, g_{n4}]$ for each $\gamma \in \pi_1(M)$ such that $I_\gamma(\rho) = \text{tr } \rho(\gamma)$ for each $\rho \in \mathfrak{R}(M)$.

We identify $\mathcal{M}_i$ and l_i with generators of $\text{im}(\pi_1(T_i) \rightarrow \pi_1(M))$. Thus, $\mathcal{M}_i$ and l_i are words in the generators for $\pi_1(M)$, and we define the following polynomial equations in the ring $\mathbb{C}[g_{11}, \dots, g_{n4}, m_1^{\pm 1}, l_1^{\pm 1}, \dots, m_h^{\pm 1}, l_h^{\pm 1}]$:

$$(4) \quad I_{\mathcal{M}_i} = m_i + m_i^{-1},$$

$$(5) \quad I_{l_i} = l_i + l_i^{-1},$$

$$(6) \quad I_{\mathcal{M}_i l_i} = m_i l_i + m_i^{-1} l_i^{-1},$$

for $i = 1, \dots, h$. Let $\mathfrak{R}_E(M)$ be the variety in $\mathbb{C}^{4n} \times (\mathbb{C} - \{0\})^{2h}$ defined by J together with the above equations. For each $\rho \in \mathfrak{R}(M)$, the equations (4–6) have a solution since commuting elements of $SL_2(\mathbb{C})$ always have a common invariant subspace. The natural projection $p_1 : \mathfrak{R}_E(M) \rightarrow \mathfrak{R}(M)$ is therefore onto, and p_1 is a dominating map.

If $(a_1, \dots, a_{4n}) \in \mathfrak{R}(M)$, then there is an action of $\mathbb{Z}_2^h$ on the resulting points

$$(a_1, \dots, a_{4n}, m_1, l_1, \dots, m_h, l_h) \in \mathfrak{R}_E(M)$$

by inverting both entries of a tuple (m_i, l_i) to (m_i^{-1}, l_i^{-1}) . The group $\mathbb{Z}_2^h$ acts transitively on the fibres of p_1 . The map p_1 is therefore finite-to-one with degree $\leq 2^h$. The maximal degree is in particular achieved when the interior of M admits a complete hyperbolic structure of finite volume, since Theorem 10 implies that there are points in $\mathfrak{X}_0(M) - \cup_{i=1}^h \{I_{\mathcal{M}_i}^2 = 4\}$.

The *eigenvalue variety* $\mathfrak{E}(M)$ is the closure of the image of $\mathfrak{R}_E(M)$ under projection onto the coordinates $(m_1, \dots, l_h)$. It is therefore defined by an ideal of the ring $\mathbb{C}[m_1^{\pm 1}, l_1^{\pm 1}, \dots, m_h^{\pm 1}, l_h^{\pm 1}]$ in $(\mathbb{C} - \{0\})^{2h}$.

Note that this construction factors through a variety $\mathfrak{X}_E(M)$, which is the character variety with its coordinate ring appropriately extended, since we can choose coordinates of $\mathfrak{X}(M)$ such that I_γ (as a function on $\mathfrak{X}(M)$) is an element of the coordinate ring of $\mathfrak{X}(M)$ for each $\gamma \in \pi_1(M)$. We let $t_E : \mathfrak{R}_E(M) \rightarrow \mathfrak{X}_E(M)$ be the natural quotient map, which is equal to $t : \mathfrak{R}(M) \rightarrow \mathfrak{X}(M)$ on the first $4n$ coordinates, and equal to the identity on the remaining $2h$ coordinates.

There also is a restriction map $r : \mathfrak{X}(M) \rightarrow \mathfrak{X}(T_1) \times \dots \times \mathfrak{X}(T_h)$, which arises from the inclusion homomorphisms $\pi_1(T_i) \rightarrow \pi_1(M)$, and we therefore denote the map $\mathfrak{X}_E(M) \rightarrow$

$\mathfrak{E}(M)$ by r_E . Denote the closure of the image of r by $\mathfrak{X}_\partial(M)$. There is the following commuting diagram of dominating maps:

$$\begin{array}{ccccc} \mathfrak{R}_E(M) & \xrightarrow{t_E} & \mathfrak{X}_E(M) & \xrightarrow{r_E} & \mathfrak{E}(M) \\ p_1 \downarrow & & p_2 \downarrow & & p_3 \downarrow \\ \mathfrak{R}(M) & \xrightarrow{t} & \mathfrak{X}(M) & \xrightarrow{r} & \mathfrak{X}_\partial(M) \end{array}$$

Note that the maps p_1, p_2, t and t_E have the property that every point in the closure of the image has a preimage, and that the maps p_1, p_2 and p_3 are all finite-to-one of the same degree.

Recall the construction by Culler and Shalen. We start with a curve $C \subset \mathfrak{X}(M)$ and an irreducible component R_C in $\mathfrak{R}(M)$ such that $t(R_C) = C$ to obtain the tautological representation $\mathcal{P} : \pi_1(M) \rightarrow SL_2\mathbb{C}(R_C)$. Let R'_C be an irreducible component of $\mathfrak{R}_E(M)$ with the property that $p_1(R'_C) = R_C$. Then $\mathbb{C}(R'_C)$ is a finitely generated extension of $\mathbb{C}(R_C)$, and we may think of $\mathcal{P}$ as a representation $\mathcal{P} : \pi_1(M) \rightarrow SL_2\mathbb{C}(R'_C)$.

If $\overline{r_E t_E(R'_C)}$ contains a curve E , then $\mathbb{C}(R'_C)$ is a finitely generated extension of $\mathbb{C}(E)$, and hence to each ideal point of E we can associate essential surfaces using $\mathcal{P}$. Since the eigenvalue of at least one peripheral element blows up at an ideal point of E , the associated surfaces necessarily have non-empty boundary (see [5]).

Thus, if there is a closed essential surface associated to an ideal point of C , then either $\overline{r_E t_E(R'_C)}$ is 0-dimensional, or there is a neighbourhood U of an ideal point ξ of R'_C such that there are (finite) points in $\overline{r_E t_E(U)} - r_E t_E(U)$. The later are called *holes in the eigenvalue variety*, examples of which are given in [17].

2.7. Boundary slopes. We denote the logarithmic limit set of $\mathfrak{E}(M)$ by $\mathfrak{E}_\infty(M)$. There are symmetries of the eigenvalue variety which give rise to symmetries of its logarithmic limit set. If $(m_1, \dots, m_i, l_i, \dots, l_h) \in \mathfrak{E}(M)$, so is $(m_1, \dots, m_i^{-1}, l_i^{-1}, \dots, l_h)$ for any i . Thus, if $\xi = (x_1, \dots, x_{2i-1}, x_{2i}, \dots, x_{2h}) \in \mathfrak{E}_\infty(M)$, then $(x_1, \dots, -x_{2i-1}, -x_{2i}, \dots, x_{2h}) \in \mathfrak{E}_\infty(M)$ for $i = 1, \dots, h$. If we factor by these symmetries, we obtain a quotient of the logarithmic limit set in $\mathbb{R}P^{2h-1}/\mathbb{Z}_2^{h-1} \cong S^{2h-1}$. The quotient map extends to a map $\Psi : S^{2h-1} \rightarrow S^{2h-1}$ of spheres, which has degree 2^h . Let

$$T = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

and let T_h be the block diagonal matrix with h copies of T on its diagonal. Then T_h is orthogonal, and its restriction to S^{2h-1} is a map of degree one.

Lemma 11 (Boundary slopes). *Let M be an orientable, irreducible, compact 3-manifold with non-empty boundary consisting of a disjoint union of h tori. If $\xi \in \mathfrak{E}_\infty(M)$ is a point with rational coordinate ratios, then there is an essential surface with boundary in M whose projectivised boundary curve coordinate is $\Psi(T_h\xi)$.*

Note that the composite ΨT_h is a smooth map of degree 2^h . Since the set of rational points is dense in the logarithmic limit set, the image $\Psi(T_h\mathfrak{E}_\infty(M))$ is a closed subset of $\mathfrak{BC}(M)$.

Remark. For a manifold M with boundary consisting of a single torus, Lemma 11 coincides with the result of [5] that slopes of sides of the Newton polygon of the A -polynomial are strongly detected boundary slopes of M .

In this case, we may normalise $\mathfrak{BC}(M)$ and $\mathfrak{E}_\infty(M)$ such that their elements are pairs of coprime integers. Assume that there is a side of the Newton polygon of slope $-p/q$ where $p, q \geq 0$ with outward pointing normal vector $\xi = (q, p)^T$. Then $\xi \in \mathfrak{E}_\infty(M)$ according to Section 1.3. We now perform matrix multiplication:

$$T\xi = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} q \\ p \end{pmatrix} = \begin{pmatrix} p \\ -q \end{pmatrix},$$

and arrive at a point $[p, -q] \in \mathfrak{BC}(M)$. The analysis for the remaining cases is similar, and we have recovered the known relationship.

Proof of Lemma 11. The coordinate ratios of ξ are rational, and hence Lemma 6 provides a curve C in $\mathfrak{E}(M)$ such that $\xi \in C_\infty$. From description (2) of elements in the logarithmic limit set, we know that ξ encodes the definition of a real-valued valuation on the function field of C :

$$\xi = (-v(m_1), \dots, -v(l_h)).$$

Since the coordinate ratios of ξ are rational, we may rescale the valuation by $r > 0$, such that for all i , $rv(m_i)$ and $rv(l_i)$ are integers, and such that there is $\alpha \in \mathbb{Z}^{2h} - \{0\}$ with $rv(m_1^{\alpha_1} \cdots l_h^{\alpha_{2h}}) = 1$. Thus, rv is a normalised, discrete, rank 1 valuation of $\mathbb{C}(C)$. We finish the proof with standard arguments. For details see Section 2.1 of [12], the proof of Proposition 2.3 of [9] or Section 5.6 of [5].

The construction by Culler and Shalen yields an action of $\pi_1(M)$ on a simplicial tree $\mathcal{T}$ associated to a finitely generated extension F of $\mathbb{C}(C)$ and to a valuation $w = dv$ for some $d > 0$. The translation length of $\gamma \in \pi_1(M)$ with eigenvalue e (as an element of F)

is $\ell(\gamma) = |2w(e)|$ (see [15], Section 3.9). A dual essential surface S associated to the action can be chosen such that the geometric intersection number of a loop in M representing γ with S is equal to $\ell(\gamma)$ for all $\gamma \in \text{im}(\pi_1(T_i) \rightarrow \pi_1(M))$. This implies that a boundary curve of S on T_i is homotopic to one of $p_i\mathcal{M}_i \pm q_i l_i$, where $n_i p_i = |2w(l_i)|$ and $n_i q_i = |2w(m_i)|$ and p_i, q_i are coprime. We need to determine the relative sign.

If $p_i\mathcal{M}_i + q_i l_i$ represents an element of the subgroup $\text{im}(\pi_1(\partial S) \rightarrow \pi_1(M))$, it is contained in the stabilizer of an edge, and hence has translation length equal to 0 since the action is without inversions. The elements $\mathcal{M}_i$ and $\mathcal{M}_i^{-1}$ translate points in opposite directions along their axis. Since l_i commutes with $\mathcal{M}_i$, they have a common axis. Thus, the curve $p_i\mathcal{M}_i + q_i l_i$ can have zero translation length only if $\mathcal{M}_i$ and l_i translate in opposite directions. This is the case if and only if $w(m_i)$ and $w(l_i)$ have opposite signs. Thus, the relative sign of p_i and q_i is opposite the relative sign of $w(m_i)$ and $w(l_i)$. We conclude that a boundary curve of S on T_i corresponds to the unoriented homotopy class of the element $\frac{1}{n_i}(2w(l_i)\mathcal{M}_i - 2w(m_i)l_i)$. Thus, the boundary curves of S are encoded by:

$$\begin{aligned} & (2w(l_1), -2w(m_1), \dots, 2w(l_h), -2w(m_h)) \\ &= 2d(v(l_1), -v(m_1), \dots, v(l_h), -v(m_h)) \\ &= 2d(T_h \xi) \end{aligned}$$

This completes the proof. ■

2.8. Dimension of the eigenvalue and character varieties. We now deduce some bounds on the dimension of the eigenvalue and the character variety.

Proposition 12. *Let M be an orientable, irreducible, compact 3-manifold with boundary consisting of h tori. Then $\dim_{\mathfrak{C}} \mathfrak{E}(M) \leq h$.*

Moreover, if the interior of M admits a complete hyperbolic structure of finite volume, then $\dim_{\mathfrak{C}} \mathfrak{E}(M) = h$.

Proof. We have $h > \dim_{\mathbb{R}} \mathfrak{BC}(M) \geq \dim_{\mathbb{R}} \mathfrak{E}_{\infty} = \dim_{\mathfrak{C}} \mathfrak{E}(M) - 1$, where the first inequality is due to Theorem 9, the second to Lemma 11, and the equality to Theorem 5. It follows that $h \geq \dim_{\mathfrak{C}} \mathfrak{E}(M)$.

To prove the second statement, it is enough to show that under the assumptions, there is a subvariety of $\mathfrak{E}(M)$ of dimension $\geq h$. Theorem 10 asserts that there is a component $\mathfrak{X}_0(M)$ of $\mathfrak{X}(M)$ of complex dimension equal to h . Thurston shows in [16], Section 5.8,

that the map $\mathfrak{X}_0(M) \rightarrow \mathbb{C}^h$ defined by $\chi_\rho \rightarrow (I_{\mathcal{M}_1}(\rho), \dots, I_{\mathcal{M}_h}(\rho))$ maps a small open neighbourhood of χ_0 to an open neighbourhood of $(\pm 2, \dots, \pm 2)$. This is only possible if $I_{\mathcal{M}_i} = \text{tr } \rho(\mathcal{M}_i)$ are algebraically independent over $\mathbb{C}$ as elements of $\mathbb{C}[X_0]$. There is a component X'_0 of $\mathfrak{X}_E(M)$ with the same properties, since p_2 is a finite dominating map and the induced homomorphism $p_{2*} : \mathbb{C}[X_0] \rightarrow \mathbb{C}[X'_0]$ is injective.

Let $E_0 = \overline{r_E t_E(X'_0)}$. If the functions $m_i + m_i^{-1}$, $i = 1, \dots, h$, are algebraically dependent over $\mathbb{C}$ as elements of $\mathbb{C}[E_0]$, then they are also algebraically dependent over $\mathbb{C}$ as elements of $\mathbb{C}[X'_0]$ since we have a ring homomorphism $\mathbb{C}[E_0] \rightarrow \mathbb{C}[X'_0]$. But the identities (4) imply that then the functions $I_{\mathcal{M}_i}$ are algebraically dependent over $\mathbb{C}$ as elements of $\mathbb{C}[X'_0]$, contradicting the fact that p_{2*} is injective.

Thus, the functions $m_i + m_i^{-1}$, $i = 1, \dots, h$, are algebraically independent over $\mathbb{C}$ as elements of $\mathbb{C}[E_0]$, and hence $\dim_{\mathbb{C}} E_0 \geq h$ according to ([7], 9.5 Theorem 2). This completes the proof. $\blacksquare$

Corollary 13. *Let M be an orientable, irreducible, compact 3-manifold with boundary consisting of h tori.*

If M contains no closed essential surface, then $\dim_{\mathbb{C}} \mathfrak{X}(M) = \dim_{\mathbb{C}} \mathfrak{E}(M) \leq h$.

Proof. We have $\dim_{\mathbb{C}} \mathfrak{X}(M) = \dim_{\mathbb{C}} \mathfrak{X}_E(M)$. Assume there is a component X of $\mathfrak{X}_E(M)$ such that $\dim X > \dim \overline{r_E(X)}$. Then [13], Theorem (3.13), implies that there is a subvariety V of dimension ≥ 1 in X which maps to a single point of $\mathfrak{E}(M)$. Associated to an ideal point of a curve in V , we find a closed essential surface since the traces of all peripheral elements are constant on V . This contradicts our assumption on M . Thus, for all components X of $\mathfrak{X}(M)$, we have $\dim X = \dim \overline{r_E(X)}$, and this proves the claim. $\blacksquare$

2.9. Remark. Let M be an orientable, irreducible, compact 3-manifold with boundary consisting of h tori, such that the interior of M admits a complete hyperbolic structure of finite volume. The proof of Proposition 12 implies that the restriction of r to the Dehn surgery component, $r_0 : X_0 \rightarrow \mathfrak{X}_\partial(M)$, is finite-to-one onto its image. The degree of r_0 depends upon $H^1(M, \mathbb{Z}_2)$, so let us consider the corresponding map $\bar{r}_0 : \bar{X}_0 \rightarrow \bar{\mathfrak{X}}_\partial(M)$. In the case where $h = 1$, Dunfield has shown in [9] that $\bar{r}_0$ has degree 1 onto its image using Thurston's Hyperbolic Dehn Surgery Theorem and a Volume Rigidity Theorem (Gromov–Thurston–Goldman). It would be interesting to generalise Dunfield's proof to arbitrary h , showing that $\bar{r}_0$ is always a birational isomorphism onto its image.

2.10. $PSL_2(\mathbb{C})$ -eigenvalue variety. Analogous to $\mathfrak{E}(M)$, one can define a $PSL_2(\mathbb{C})$ -eigenvalue variety $\overline{\mathfrak{E}}(M)$, since the function $I_\gamma^2 : \overline{\mathfrak{X}}(M) \rightarrow \mathbb{C}$ defined by $I_\gamma^2(\bar{\rho}) = (\text{tr } \bar{\rho}(\gamma))^2$ is regular (i.e. polynomial) for all $\gamma \in \pi_1(M)$ (see [3]). Thus, we construct a variety $\overline{\mathfrak{R}}_E(M)$ using the relations

$$\begin{aligned} I_{\mathcal{M}_i}^2 &= M_i + 2 + M_i^{-1}, \\ I_{l_i}^2 &= L_i + 2 + L_i^{-1}, \\ I_{\mathcal{M}_i l_i}^2 &= M_i L_i + 2 + M_i^{-1} L_i^{-1}, \end{aligned}$$

for $i = 1, \dots, h$.

Consider the representation of $\mathbb{Z} \oplus \mathbb{Z}$ into $PSL_2(\mathbb{C})$ generated by the images of

$$(7) \quad \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

In $PSL_2(\mathbb{C})$, the image of this representation is isomorphic to $\mathbb{Z}_2 \oplus \mathbb{Z}_2$, but the image of any lift to $SL_2(\mathbb{C})$ is isomorphic to the quaternion group Q_8 (in Quaternion group notation). If $\bar{\rho} \in \overline{\mathfrak{R}}(M)$ restricts to such an irreducible abelian representation on a boundary torus T_i , then the equations $M_i + 2 + M_i^{-1} = 0$, $L_i + 2 + L_i^{-1} = 0$, $M_i L_i + 2 + M_i^{-1} L_i^{-1} = 0$ have no solution. In particular, the projection $\bar{\rho} : \overline{\mathfrak{R}}_E(M) \rightarrow \overline{\mathfrak{R}}(M)$ may not be onto.

The closure of the image of $\overline{\mathfrak{R}}_E(M)$ onto the coordinates $(M_1, L_1, \dots, M_h, L_h)$ is denoted by $\overline{\mathfrak{E}}(M)$ and called the $PSL_2(\mathbb{C})$ -eigenvalue variety. The relationship between ideal points of $\overline{\mathfrak{E}}(M)$ and strongly detected boundary curves is the same as for the $SL_2(\mathbb{C})$ -version, since the proof of Lemma 11 applies without change:

Lemma 14. *Let M be an orientable, irreducible, compact 3-manifold with non-empty boundary consisting of a disjoint union of h tori. If $\xi \in \overline{\mathfrak{E}}_\infty(M)$ is a point with rational coordinate ratios, then there is an essential surface with boundary in M whose projectivised boundary curve coordinate is $\Psi(T_h \xi)$.*

3. TORUS KNOTS

In general, one needs a computer to calculate the A -polynomial. It is shown in [5] that if $\mathfrak{k}$ is a nontrivial (p, q) -torus knot, then $A_{\mathfrak{k}}(l, m)$ is divisible by $lm^{pq} + 1$ (where we follow the convention that if M is the complement of a knot $\mathfrak{k}$ in S^3 , then $\{\mathcal{M}, l\}$ is a standard peripheral system).

Proposition 15. ¹ *Let $\mathfrak{k}$ be a (p, q) -torus knot. If $p = 2$ or $q = 2$, then $A_{\mathfrak{k}}(l, m) = (l - 1)(lm^{pq} + 1)$, otherwise $A_{\mathfrak{k}}(l, m) = (l - 1)(lm^{pq} + 1)(lm^{pq} - 1)$.*

Proof. The fundamental group of $S^3 - \mathfrak{k}$ is presented by $\Gamma = \langle u, v \mid u^p = v^q \rangle$, and a standard peripheral system is given by $\mathcal{M} = u^n v^m$ and $l = u^p \mathcal{M}^{-pq}$, where $mp + nq = 1$. These facts can be found in [4] on page 45. The factor $l - 1$ arises from reducible representations, and all factors arising from components containing irreducible representations are to be determined.

The element u^p is in the centre of Γ , since it is identical to v^q and hence commutes with both generators. Thus the image of u^p is in the center of $\rho(\Gamma)$. Since two commuting elements have a common eigenvector, $\rho(u)^p$ has a common eigenvector with $\rho(u)$ and $\rho(v)$ respectively. If the representation is irreducible, these eigenvectors have to be distinct. Thus, after conjugation it may be assumed that the generators map to upper and lower triangular matrices respectively, and the central element is represented by a diagonal matrix. Direct matrix computations show that the commutativity with either of the generators gives $\rho(u)^p = \pm E$.

Assume that $\rho(u)^p = -E$. The relation $\rho(l) = -\rho(\mathcal{M})^{-pq}$, which is equivalent to $\rho(l\mathcal{M}^{pq}) = -E$, then implies the equation $lm^{pq} = -1$. This is the curve obtained in [5] by sending u and v to noncommuting elements of $SL_2(\mathbb{C})$ of order exactly $2p$ and $2q$ respectively.

If the image of $\rho(u)^p$ is trivial, then $\rho(v)^q = \rho(u)^p = E$. If p or q equals 2, this implies that the image of one of the generators is $\pm E$. But this yields that $\rho(\Gamma)$ is abelian, and therefore contradicts the irreducibility assumption. This completes the proof of the first assertion.

Now assume that neither of p and q is equal to two. As above, curves of irreducible representations are obtained by sending u and v to noncommuting elements of $SL_2(\mathbb{C})$ of order exactly p and q respectively. Any of these curves yields a component of the eigenvalue variety defined by the equation $lm^{pq} = 1$, and this finishes the proof of the proposition. In fact, with a more precise analysis one could count the number of 1-dimensional curves in the character variety of a torus knot. ■

¹I thank Walter Neumann for pointing out that an earlier argument for $(\alpha, 1)$ -two bridge knots could be generalised to torus knots.

Similarly to the above, we can compute a generator for the defining (principal) ideal of the $PSL_2(\mathbb{C})$ -eigenvalue variety of a torus knot. We call this generator the $\overline{A}$ -polynomial, and its variables have the unfortunate names L and M . Thus, $\overline{A}_M(L, M)$ is the $\overline{A}$ -polynomial associated to the manifold M .

Proposition 16. *Let $\mathfrak{k}$ be a (p, q) -torus knot. Then $\overline{A}_{\mathfrak{k}}(L, M) = (L - 1)(LM^{pq} - 1)$.*

Proof. It follows from the lemma in Section 6 of [5] that all $PSL_2(\mathbb{C})$ -representations of a knot group lift to $SL_2(\mathbb{C})$. Thus, all factors of the A -polynomial arise from factors of the $\overline{A}$ -polynomial and vice versa.

The eigenvalue of the meridian appears with even powers in the (unfactorised) A -polynomial of $\mathfrak{k}$, and the sign of the longitude's eigenvalue is uniquely determined since l is null-homologous. Thus, $A_{\mathfrak{k}}(l, m) = A_{\mathfrak{k}}(l, -m)$, but $A_{\mathfrak{k}}(l, m) = 0$ does not imply $A_{\mathfrak{k}}(-l, m) = 0$ in general. However, since $\overline{A}_{\mathfrak{k}}(l^2, m^2) = \overline{A}_{\mathfrak{k}}((-l)^2, m^2)$, it follows that the variety defined by $\overline{A}_{\mathfrak{k}}(l^2, m^2) = 0$ is the locus of $A_{\mathfrak{k}}(l, m)A_{\mathfrak{k}}(-l, m) = 0$. Expanding $A_{\mathfrak{k}}(l, m)A_{\mathfrak{k}}(-l, m)$, substituting $l^2 = L$ and $m^2 = M$ and deleting repeated factors therefore gives the result as stated. ■

Acknowledgements. The contents of this paper forms part of my Ph.D. thesis, and I thank my supervisors Craig Hodgson and Walter Neumann for their constant support and inspiration. This research was supported in part by an International Postgraduate Research Scholarship by the Commonwealth of Australia Department of Education, Science and Training.

REFERENCES

- [1] G.M. Bergman: *The logarithmic limit-set of an algebraic variety*, Trans. Amer. Math. Soc., 157, 459-469, (1971).
- [2] R. Bieri, J.R.J.Groves: *The geometry of the set of characters induced by valuations*, Jour. reine und angew. Math., 347, 168-195, (1984).
- [3] S. Boyer, X. Zhang: *On Culler-Shalen seminorms and Dehn filling*, Ann. Math., 148, 737-801 (1998).
- [4] G. Burde, H. Zieschang: *Knots*, Walter de Gruyter, Berlin New York, 1985.
- [5] D. Cooper, M. Culler, H. Gillett, D.D. Long and P.B. Shalen: *Plane curves associated to character varieties of 3-manifolds*, Invent. Math., 118, 47-84 (1994).
- [6] D. Cooper, D.D. Long: *Representation Theory and the A-polynomial of a knot*, Chaos, Solitons & Fractals, 9, 749-763 (1998).
- [7] D. Cox, J. Little, D. O'Shea: *Ideals, Varieties, and Algorithms*, Springer, New York, 1992.

- [8] M. Culler, P.B. Shalen: *Varieties of group representations and splittings of 3-manifolds*, Ann. Math., 117, 109-146 (1983).
- [9] N.M. Dunfield: *Cyclic Surgery, degrees of maps of character curves, and volume rigidity for hyperbolic manifolds*, Invent. Math., 136, 623-657 (1999).
- [10] A.E. Hatcher: *On the boundary curves of incompressible surfaces*, Pacific Jour. Math., 99, 373-377, (1982).
- [11] A.E. Hatcher: *Notes on Basic 3-Manifold Topology*, available from <http://math.cornell.edu/~hatcher>.
- [12] A.E. Lash: *Boundary Curve Space of the Whitehead Link Complement*, Ph. D. thesis, UC Santa Barbara, 1993.
- [13] D. Mumford: *Algebraic Geometry I Complex Projective Varieties*, Springer, Berlin, 1976.
- [14] I.R. Shafarevich: *Basic Algebraic Geometry 1 & 2*, Springer Verlag, Berlin, 1994.
- [15] P.B. Shalen: *Representations of 3-manifold groups*, Handbook of geometric topology, 955-1044, Amsterdam, 2002.
- [16] W.P. Thurston: *The geometry and topology of 3-manifolds*, Princeton Univ. Math. Dept. (1978). Available from <http://msri.org/publications/books/gt3m/>.
- [17] S. Tillmann: *On the Kinoshita-Terasaka mutants and generalised Conway mutation*, J. Knot Theory Ramifications, 9, 557-575 (2000).
- [18] S. Tillmann: *On character varieties*, Ph.D. thesis, The University of Melbourne, 2002.

Department of Mathematics and Statistics, The University of Melbourne, VIC 3010, Australia